

Yoga For The Adjunctive Management Of Chronic Low Back Pain: A Clinical Review

Neil K. Agarwal (MD)¹ & Shashi K. Agarwal (MD)²

1. Internal Medicine Resident, Hahnemann University Hospital, 230 North Broad Street, Philadelphia, PA 19102, USA, Email: neil.k.agarwal@gmail.com
2. Center for Contemporary and Complimentary Cardiology, 62, Chester Circle, #2B, New Brunswick, NJ, USA, Email: usacardiologist@gmail.com

ABSTRACT

Introduction: Chronic low back pain is a major cause of activity limitation and work absence worldwide. Despite a wide range of conventional pharmacologic agents, non-pharmacologic interventions, and surgical procedures, most patients report only mild or moderate relief. It continues to exert an enormous social and economic burden on individuals, families, communities, industry and the governments. Yoga is not only efficacious, but easy to practice, safe and cost effective in ameliorating this chronic malady.

Methods: A systemic review of scientific publications cited by PubMed was done. There were 3857 citations under ‘yoga’ dating back to 1948 and 8992 citations under ‘chronic low back pain’ dating back to 1946. There were 92 entries under ‘yoga and chronic low back pain’, dating back to 1982. Other relevant published scientific material was also consulted.

Results: Several large and well-designed studies support yoga’s effectiveness for reducing pain, increasing spinal flexibility and improving functionality in patients with chronic low back pain. Pain medication usage is reduced, fatigue is diminished and there is an amelioration of depression and anxiety. Yoga appears to be more effective than physical therapy, manipulative procedures and other complementary modalities like acupuncture, in this condition. Compliance is good and the benefits are long lasting.

Conclusion: Low back pain is associated with a substantial morbidity, disability and costs to the Indian society. Yoga is backed by evidence based data supporting its efficacy and safety as an adjunctive therapy in the management of chronic low back pain.

Introduction

Chronic low back pain is defined as low back pain of at least three month’s duration. (Croft et al., 1998) It is among the most common of all health complaints. (Rives et al, 2004) Yearly prevalence ranges from 22% to 65% and lifetime prevalence from 11% to 84%. (Walker, 2000) It is a major cause of activity limitation and work absence worldwide. (Hoy et al, 2012) The Global Burden of Disease 2010 estimates that low back pain is amongst the top ten DALYs (disability-adjusted life years) causing diseases and injuries in the world. (Murray et al, 2012) In the developed countries like the United States, yearly prevalence of back pain varies from 10% to 56% (Deyo et al, 1987)

with a lifetime prevalence of 65% to 80% (Lawrence et al, 1998) It is considered the most expensive benign medical condition in the industrialized world. (Engel et al, 1996)

Its yearly prevalence in the developing countries is not much lower, varying between 36% and 64%. (Hoy et al., 2003; Cakmak et al, 2004; Gilgil et al, 2005; Barrero et al, 2006) South Asia is no exception. (Bindra et al, 2015) Its prevalence in the Indian population varies between 6.2% in the general population to 92% in construction workers. (Bindra et al, 2015) Etiological factors may like tuberculosis are unique to the Indian population. (Krishna et al, 2014) The patients with chronic low back pain in South Asia are usually from the lower socio-economic status,

do not visit a consultant, and often prefer over-the-counter medications or traditional treatments. (Bindra et al, 2015) Chronic low back pain is also common in the poorer countries of Africa. (Omokhodion et al, 2003; Omokhodion 2004; Gilgil et al, 2005) In a systemic review, the average lifetime prevalence of low back pain in Africa, among the adolescents was 36% and among adults was 62%. (Louw et al. 2007)

Despite a wide range of conventional pharmacologic agents, non-pharmacologic interventions, and surgical procedures, most patients report only mild or moderate relief. (van Tulder et al, 2000; Balague et al, 2012; Haldeman et al, 2008; Bogduk, 2004; Wolsko et al, 2003) Prognosis remains poor. (Carey et al, 2000; Grotle et al, 2006) Chronic low back pain continues to exert an enormous social and economic burden on individuals, families, communities, industry and the state governments. (Woolf et al, 2003; Katz, 2006) Yoga provides a bio-psycho-social rehabilitation that is successful in the management of this condition. (Guzman et al, 2002) It is easy to learn and practice, relatively safe, and cheap. (Holtzman et al, 2013) This brief review focusses on the increasing scientific evidence confirming the beneficial effects of yoga as an adjunctive therapeutic modality in patients with chronic low back pain.

Methods

A comprehensive literature search was carried out using the PubMed database of the US National Library of Medicine, National Institutes of Health in February of 2017. Other contemporary and complementary medicine databases (including PMC, Medline, Google Scholar, and Quertile) were also queried and relevant publications were consulted. Other available and relevant published scientific material, identified through references, was also used. Only English language literature was reviewed.

Results

There were 3857 citations under 'yoga' dating back to 1948 and 8992 citations under

'chronic low back pain' dating back to 1946. There were 92 entries under 'yoga and chronic low back pain', dating back to 1982. There were 539 citations under 'yoga and pain' dating back to 1963; 102 under 'yoga and muscle strength' dating back to 1991; 121 under 'yoga and flexibility' dating back to 1964; 149 under 'yoga and balance' dating back to 1975; 494 citations under 'yoga and depression' dating back to 1974; 522 under 'yoga and anxiety' dating back to 1968; 80 under 'yoga and sleep disorders' dating back to 1963 and 606 under 'yoga and quality of life' dating back to 1987.

Discussion

Several large and well-designed studies support yoga's effectiveness in patients with chronic low back pain. (Galantino et al, 2004, Jacobs et al, 2004, Williams et al, 2005, Sherman et al, 2005, Holtzman et al, 2013) The benefits of yoga stem from this biopsychosocial modality's (Gatchel et al, 2007) ability to beneficially ameliorate the litany of physical and emotional limitations encountered by these patients. There are measurable modulations of several neuroendocrine mediators, with beneficial results. (Sherman et al, 2013; Lee et al, 2014) Yoga appears to be more effective than physical therapy, manipulative procedures and other complementary modalities like acupuncture, in treating this condition. It is extremely cost-effective. Compliance is good, (Caren et al, 2015) and the benefits are long lasting.

Pain: Pain is the main disabling symptom, experienced by 80%-90% patients with chronic back discomfort. (Kovacs et al, 2005) The aim of therapeutic modalities is thus aimed primarily at reducing pain, preventing pain catastrophizing and increasing pain tolerance. Pain medications, both nonsteroidal anti-inflammatory drugs and opiates are commonly used as analgesics. (Moulin et al, 1996; van Tulder et al, 2000) However, relief is usually only temporary, and there is no associated psychological benefit. Anti-depressants (Salerno et al, 2002) and muscle

relaxants also provide only limited relief. (van Tulder et al, 2003) Many other therapies, such as acupuncture and massage are only mildly more effective than sham therapy and invariably provide only temporary relief. They have limited success in long term improvements. (Cherkin et al, 2003; McIlveen et al, 1998) Surgical interventions provide some relief but are invasive and often non-lasting. (Fritzell et al, 2001) Many other manipulative treatments have been tried without any persuasive efficacy. (Furlan et al, 2010) Exercise is a major non-pharmacological intervention, (O'Sullivan et al, 1997; Hayden et al, 2005) and is commonly utilized in the rehabilitation processes. However, results are less than impressive. Yoga has emerged as the forerunner in the treatment of pain associated with the low back. (Evans et al, 2010) It favorably affects all parameters of pain. (Carson et al, 2010; Curtis et al, 2011; John et al, 2007; Tilbrook et al, 2011) Since pain has a strong 'mind-connection' component, (Kabat-Zinn et al, 1986) yoga by "uncoupling" the physical sensation, from the emotional and cognitive experience of pain, contributes to the relief process. The pain reduction persists for several months during the post-treatment follow up period, (Jacobs et al, 2004; Sherman et al, 2005) and pain medication usage is considerably reduced.

Muscle strength and endurance, joint flexibility, posture and balance: Yoga uses the body's own weight and earth's gravity to put the various parts of the body through a range of motion. These postures gradually increase muscle strength and joint flexibility, (Carneiro et al, 2010; Field, 2011, Ryba et al, 2006; Roland et al, 2011; Caren et al, 2015) both in healthy individuals (Tran et al, 2001) as well as in patients with chronic low back pain. (Tekur et al, 2008) Muscular endurance is improved. (Chen et al, 2008) Yoga postures put the spine through a wide range of motions and help improve spinal alignment and bodily posture. (Gail et al, 2009) Balance is improved and falls are reduced, especially in the elderly. (Saravanakumar et al, 2014; Vaughan et al,

2014) These musculoskeletal improvements help ameliorate chronic low back pain.

Psychological factors: Associated psychological impairments are extremely common in patients with chronic low back pain. (Demyttenaere et al, 2007; Linton, 2000) The three common comorbidities are depression, anxiety and sleep disorders. (Patten, 2001; Newcomer et al, 2010; Bahouq et al, 2013) Depression incidence is doubled in patients with chronic low back pain, (Patten, 2001) and appears to have a bidirectional association: depression is a predictor of persistent pain and pain is a predictor of persistence of depression. (Saito et al, 2012; Ohayon et al, 2003)) There is strong evidence from several randomized trials supporting the benefits of yoga in reducing depression. (Groessler et al, 2008; Kinser et al, 2012) Depression in patients with low back pain prognosticate a more refractory and longer therapeutic course and more workplace time lost. (Bair et al, 2003) The economic burden is much higher on the society in these patients. (Gore et al, 2012) The beneficial role of yoga in alleviating anxiety is also well documented. (Hofman et al, 2015) Sleep disorders are also a common co-morbidity of chronic back pain, (Bahouq et al, 2013, Kathi et al, 2012) and yoga helps in improving sleep in these patients. (Balasubramaniam et al, 2012) Yoga increases functionality in patients with chronic back pain. (Telles et al, 2011; Holtzman et al, 2013) There are documented improvements in the quality of life in these patients. (Oken et al, 2006; Williams et al, 2009; Tekur et al, 2010; Banth et al, 2015) Patients with chronic low back pain also suffer from strained interpersonal relationships and financial difficulties. (Anderson, 1999; Balaque et al, 2012; Linton, 2000) There is also the positive effect provided by a sense of belonging when attending a yoga class – resulting in an increase in emotional and tangible support. (Wren et al, 2011) Yoga therefore not only helps ameliorate chronic low back pain but also helps rectify its associated psychological comorbidities.

Conclusion

Low back pain is associated with a substantial morbidity, disability and costs all over the world. In developed countries like the USA, causes include work related injuries, prolonged sitting, motor vehicle accidents and falls. In developing countries like India, labor jobs often involve improperly lifting of heavy loads. People of lower socioeconomic status also have poor access to consultants, end up taking over the counter pain medications and

may prefer traditional treatments. Yoga is backed by evidence based data supporting its efficacy and safety as an adjunctive therapy in the management of chronic low back pain. Yoga is increasingly popular all over the world, easy to learn and perform and basically cheap or even free. It is therefore ideally suited as an adjunctive therapy for patients with chronic low back pain.

Conflict of interest: none

Acknowledgements: none

References:

1. Andersson GBJ. Epidemiological features of chronic low-back pain. *Lancet*. 1999; 354:581–5.
2. Bahouq H, Allali F, Rkain H, et al. Prevalence and severity of insomnia in chronic low back pain patients. *Rheumatol Int*. 2013; 33:1277–81.
3. Bair MJ1, Robinson RL, Katon W, et al. Depression and pain comorbidity: a literature review. *Arch Intern Med*. 2003 Nov 10;163(20):2433-45.
4. Balague F, Mannion AF, Felleise F, et al. Non-specific low back pain. *Lancet*. 2012; 379:482–91.
5. Balasubramaniam M, Shirley Telles, P. MuraliDoraiswamy. Yoga on Our Minds: A Systematic Review of Yoga for Neuropsychiatric Disorders. *Front Psychiatry*. 2012; 3: 117.
6. Banth S, Maryam Didehdar Ardebil. Effectiveness of mindfulness meditation on pain and quality of life of patients with chronic low back pain. *Int J Yoga*. 2015 Jul-Dec;8(2):128-33.
7. Barrero LH, Hsu YH, Terwedow H, et al. (2006). Prevalence and physical determinants of low back pain in a rural Chinese population. *Spine* 31 2728–2734.
8. Bindra S., Sinha A.G.K., Benjamin A.I. Epidemiology of low back pain in Indian population: A review. *International Journal of Basic and Applied Medical Sciences* 2015 Vol. 5 (1) January-April, pp. 166-179.
9. Bogduk N. Management of chronic low back pain. *Med J Aust*. 2004; 180:79-83.
10. Cakmak A, Yucel B, Ozalcin SN, et al. (2004). The frequency and associated factors of low back pain among a younger population in Turkey. *Spine* 29: 1567–1572. 2004.
11. Caren Lau, Ruby Yu, Jean Woo. Effects of a 12-Week Hatha Yoga Intervention on Cardiorespiratory Endurance, Muscular Strength and Endurance, and Flexibility in Hong Kong Chinese Adults: A Controlled Clinical Trial. *Evid Based Complement Alternat Med*. 2015; 2015: 958727.
12. Carey TS, Garrett JM, Jackman AM. Beyond the good prognosis: examination of an inception cohort of patients with chronic low back pain. *Spine* 2000; 25:115-20.
13. Carneiro KA, Rittenberg JD. The role of exercise and alternative treatments for low back pain. *Phys Med Rehabil Clin North Am*. 2010; 21:777–92.
14. Carson JW, Carson KM, Jones KD, et al. A pilot randomized controlled trial of the Yoga of Awareness program in the management of fibromyalgia. *Pain*. 2010; 151:530–9
15. Chen K. M., Chen M. H., Hong S. M., et al. Physical fitness of older adults in senior activity centres after 24-week silver yoga exercises. *Journal of Clinical Nursing*. 2008;17(19):2634–2646.
16. Cherkin DC, Sherman KJ, Deyo RA, et al. A review of the evidence for the effectiveness, safety, and cost of acupuncture, massage therapy, and spinal manipulation for back pain. *Ann Intern Med* 2003; 138: 898-906.

17. Croft PR, Macfarlane GJ, Papageorgiou AC, et al. Outcome of low back pain in general practice: a prospective study. *BMJ* 1998; 316:1356–1359.
18. Curtis K, Osadchuk A, Katz J. An eight-week yoga intervention is associated with improvements in pain, psychological functioning and mindfulness, and changes in cortisol levels in women with fibromyalgia. *J Pain Res.* 2011; 4:189–201.
19. Demyttenaere K, Bruffaerts R, Lee S, et al. Mental disorders among persons with chronic back or neck pain: results from the World Mental Health Surveys. *Pain.* 2007 Jun;129(3):332-42.
20. Deyo RA, Tsui-Wu YJ. Descriptive epidemiology of low back pain and its related medical care in the United States. *Spine* 1987; 12:264-268.
21. Engel CC, von Korff M, Katon WJ. Back pain in primary care: predictors of high health-care costs. *Pain.* 1996 May-Jun. 65(2-3):197-204.
22. Evans DD, Carter M, Panico R, et al. Characteristics and predictors of short-term outcomes in individuals self-selecting yoga or physical therapy for treatment of chronic low back pain. *PM R.* 2010; 2:1006–1015.
23. Field T. Yoga clinical research review. *Complement TherClinPract.* 2011; 17:1–8.
24. Fritzell P, Hagg O, Wessberg P, et al. Swedish Lumbar Spine Study Group. 2001 Volvo award winner in clinical studies: lumbar fusion versus nonsurgical treatment for chronic low back pain. A multicenter randomized controlled trial from the Swedish Lumbar Spine Study Group. *Spine* 2001; 26: 2521-2534.
25. Furlan AD, FatemehYazdi, Alexander Tsertsvadze, et al. Complementary and Alternative Therapies for Back Pain II. University of Ottawa Evidence-based Practice Center, Rockville (MD): Agency for Healthcare Research and Quality (US); 2010 Oct. Report No.: 10(11)-E007
26. Gail A. Greendale, Mei-Hua Huang, Arun S. Karlamangla, et al. Yoga decreases kyphosis in senior women and men with adult onset hyperkyphosis: results of a randomized controlled trial. *J Am Geriatr Soc.* 2009 Sep; 57(9): 1569–1579.
27. Galantino ML, Bzdewka TM, Eissler-Russo JL, et al. The impact of modified Hatha yoga on chronic low back pain: a pilot study. *Altern Ther Health Med.* 2004; 10:56-9.
28. Gatchel RJ, Peng YB, Peters ML, et al. The biopsychosocial approach to chronic pain: Scientific advances and future directions. *Psychol Bull.* 2007; 133:581–624.
29. Gilgil E, Kacar C, Butun B, et al. (2005). Prevalence of low back pain in a developing urban setting. *Spine* 30: 1093–1098. 2005.
30. Gore M, Sadosky A, Stacey BR, et al. The burden of chronic low back pain: clinical comorbidities, treatment patterns, and health care costs in usual care settings. *Spine* 2012 May 15; 37(11).
31. Groessl EJ, Weingart KR, Aschbacher K, et al. Yoga for veterans with chronic low-back pain. *J Altern Complement Med.* 2008;14:1123–1129.
32. Grotle M, Vollestad NK, Brox JI. Clinical course and impact of fear-avoidance beliefs in low back pain: prospective cohort study of acute and chronic low back pain II. *Spine* 2006; 31:1038-46.
33. Guzman J, Esmail R, Karjalainen K, et al. Multidisciplinary bio-psycho-social rehabilitation for chronic low back pain. *Cochrane Database Syst Rev.* 2002.
34. Haldeman S, Dagenais S. A supermarket approach to the evidence-informed management of chronic low back pain. *Spine J.* 2008; 8:1–7.
35. Hayden JA, van Tulder MW, Malmivaara AV, et al. Meta-analysis: exercise therapy for nonspecific low back pain. *Ann Intern Med.* 2005 May 3;142(9):765-75.
36. Hofmann SG, Curtiss J, Khalsa SB, et al. Yoga for generalized anxiety disorder: design of a randomized controlled clinical trial. *ContempClin Trials.* 2015 Aug 6; 44:70-76.

37. Holtzman S, Beggs RT. Yoga for chronic low back pain: A meta-analysis of randomized controlled trials. *Pain Res Manag.* 2013 Sep-Oct; 18(5): 267–272.
38. Hoy D, Toole MJ, Morgan D et al. (2003). Low back in rural Tibet. *Lancet* 361 225–226.
39. Hoy D, Christopher Bain, Gail Williams, et al. A systematic review of the global prevalence of low back pain. *Arthritis & Rheumatism*, 2012; 64(6) :2028–2037.
40. Jacobs BP, Mehling W, Avins AL, et al. Feasibility of conducting a clinical trial on Hatha yoga for chronic low back pain: methodological lessons. *Altern Ther Health Med.* 2004; 10:80-3.
41. John PJ, Sharma N, Sharma CM, et al. Effectiveness of yoga therapy in the treatment of migraine without aura: A randomized controlled trial. *Headache.* 2007; 47:654–61.
42. Kabat-Zinn J, Lipworth L, Burney R, et al. Four-Year Follow-up of a meditation-based program for the self-regulation of chronic pain: Treatment outcomes and compliance. *Clin J Pain.* 1986; 2:159–73.
43. Kathi L. Heffner, Christopher R. France, Zina Trost, et al. Chronic Low Back Pain, Sleep Disturbance, and Interleukin-6. *Clin J Pain.* 2011 Jan; 27(1): 35–41.
44. Katz JN. Lumbar disc disorders and low-back pain: socioeconomic factors and consequences. *J Bone Joint Surg Am*, 2006, 88(suppl 2):21-24.
45. Kinser PA, Lisa Goehler, Ann Gill Taylor. How Might Yoga Help Depression? A Neurobiological Perspective. *Explore (NY).* 2012 Mar 1; 8(2): 118–126.
46. Kovacs FM, Abaira V, Zamora J, et al. Spanish Back Pain Research Network. The transition from acute to subacute and chronic low back pain: A study based on determinants of quality of life and prediction of chronic disability. *Spine (Phila Pa 1976)* 2005; 30:1786–92.
47. Krishna V K, D. Sharma, G. Samuel. Epidemiological Study for Evaluation of Etiology and Risk Factors in Patients with Low Back Pain. *Global Spine J* 2014; 04 – or 1.01.
48. Lawrence RC, Helmick CG, Arnett FC. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis & Rheumatism* 1998; 41:778-799.
49. Lee M, Moon W, Kim J. Effect of yoga on pain, brain-derived neurotrophic factor, and serotonin in premenopausal women with chronic low back pain. *Evid Based Complement Alternat Med.* 2014;2014:203173.
50. Linton SJ. A review of psychological risk factors in back and neck pain. *Spine.* 2000; 25:1148–56.
51. Louw Q, Linzette D Morris and Karen Grimmer-Somers. The Prevalence of low back pain in Africa: a systematic review. *BMC Musculoskeletal Disorders* 2007, 8:105 doi:10.1186/1471-2474-8-105
52. McIlveen B, Robertson V. A randomized controlled study of the outcome of hydrotherapy for subjects with low back or back and leg pain. *Physiotherapy* 1998; 84: 17-26.
53. Moulin DE, Iezzi A, Amireh R, et al. Randomised trial of oral morphine for chronic non-cancer pain. *Lancet* 1996; 347: 143-147.
54. Murray CJL, Theo Vos, Rafael Lozano, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2197–223.
55. Newcomer KL, Shelerud RA, Vickers Douglas KS, et al. Anxiety levels, fear-avoidance beliefs, and disability levels at baseline and at 1 year among subjects with acute and chronic low back pain. *PM R* 2010; 2:514–20.
56. Ohayon MM, Schatzberg AF (2003) Using chronic pain to predict depressive morbidity in the general population. *Arch Gen Psychiatry* 60: 39–47.
57. Oken BS, Zajdel D, Kishiyama S, et al. Randomized, controlled, six-month trial of yoga in healthy seniors: effects on cognition and quality of life. *Altern Ther Health Med.* 2006;12(1):40–47.

58. Omokhodion F, Sanya A. Risk factors for low back pain among office workers in Ibadan, Southwest Nigeria. *Occupational Medicine*. 2003;53:287–89. doi: 10.1093/occmed/kqg063.
59. Omokhodion F. Low back pain in an urban population in Southwest Nigeria. *Tropical Doctor*. 2004;34:17–20
60. O’Sullivan PB, Twomey LT, Allison GT. Evaluation of specific stabilizing exercise in the treatment of chronic low back pain with radiologic diagnosis of spondylolysis or spondylolisthesis. *Spine* 1997; 22: 2959-2967.
61. Patten SB. Long-term medical conditions and major depression in a Canadian population study at waves 1 and 2. *J Affect Disord*. 2001; 63:35–41.
62. Rives PA, Douglass AB. Evaluation and treatment of low back pain in family practice. *J Am Board Fam Pract*. 2004;17: S23–31.
63. Ryba TV, Kaltenborn JM. The benefits of yoga for athletes: The body. *Athletic Therapy Today*. 2006; 11:32–34.
64. Roland KP, Jakobi JM, Jones GR. Does yoga engender fitness in older adults? A critical review. *J Aging Phys Act*. 2011; 19:62–79.
65. Saito T, Kai I, Takizawa A. Effects of a program to prevent social isolation on loneliness, depression, and subjective well-being of older adults: a randomized trial among older migrants in Japan. *Arch GerontolGeriatr* 2012; 55: 539–54.
66. Salerno SM, Browning R, Jackson JL. The effect of antidepressant treatment on chronic back pain: a meta-analysis. *Arch Intern Med*. 2002 Jan 14;162(1):19-24.
67. Saravanakumar P, Higgins IJ, van der Riet PJ, et al. The influence of tai chi and yoga on balance and falls in a residential care setting: A randomised controlled trial. *Contemp Nurse*. 2014;48(1):76-87.
68. Sherman KJ, Daniel C. Cherkin, Janet Erro, et al. Comparing Yoga, Exercise, and a Self-Care Book for Chronic Low Back Pain: A Randomized, Controlled Trial. *Ann Intern Med*. 2005;143(12):849-856.
69. Sherman KJ, Wellman RD, Cook AJ, et al. Mediators of yoga and stretching for chronic low back pain. *Evid Based Complement Alternat Med*. 2013;2013:130818.
70. Tekur P., Singphow C., Nagendra H. R., et al. Effect of short-term intensive yoga program on pain, functional disability and spinal flexibility in chronic low back pain: a randomized control study. *Journal of Alternative and Complementary Medicine*. 2008;14(6):637–644.
71. Tekur P, Chametcha S, Hongasandra RN, et al. Effect of yoga on quality of life of CLBP patients: A randomized control study. *Int J Yoga*. 2010;3:10–17.
72. Telles S, Kalkuni V Naveen, Vaishali Gaur, et al. Effect of one week of yoga on function and severity in rheumatoid arthritis. *BMC Res Notes*. 2011; 4: 118.
73. Tilbrook HE, Cox H, Hewitt CE, et al. Yoga for chronic low back pain. *Ann Intern Med*. 2011; 155:569–78.
74. Tran M. D., Holly R. G., Lashbrook J., Amsterdam E. A. Effects of hatha yoga practice on the health-related aspects of physical fitness. *Preventive Cardiology*. 2001;4(4):165–170.
75. van Tulder MW, Scholten RJPM, Koes BW, et al. Nonsteroidal anti-inflammatory drugs for low back pain. A systematic review within the framework of the Cochrane Collaboration Back Review Group. *Spine* 2000; 25: 2501-2513.
76. van Tulder MW, Touray T, Furlan AD, et al. Muscle relaxants for nonspecific low back pain: a systematic review within the framework of the Cochrane Collaboration. *Spine* 2003; 28: 1978-1992.
77. Vaughan P Nicholson, Mark R McKean, and Brendan J Burkett. Twelve weeks of BodyBalance® training improved balance and functional task performance in middle-aged and older adults. *ClinInterv Aging*. 2014; 9: 1895–1904.
78. Walker BF. The prevalence of low back pain: a systematic review of the literature from 1966 to 1998. *J Spinal Disord* 2000; 13:205-17.

79. Williams KA, Petronis J, Smith D, et al. Effect of Iyengar yoga therapy for chronic low back pain. *Pain*. 2005; 115:107-17.
80. Williams K, Abildso C, Steinberg L, Doyle E, Epstein B, et al. Evaluation of the effectiveness and efficacy of Iyengar yoga therapy on chronic low back pain. *Spine (Phila Pa 1976)* 2009;34:2066–2076.
81. Wolsko PM, Eisenberg DM, Davis RB, et al. Patterns and perceptions of care for treatment of back and neck pain: results of a national survey. *Spine*. 2003; 28:292-7.
82. Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. *Bull World of Health Organ*. 2003; 81:646–56.
83. Wren AA, Wright MA, Carson JW, et al. Yoga for persistent pain: New findings and directions for an ancient practice. *Pain*. 2011; 152:477–80.

